



9316-R2

FUL-O-PEP

16/6 EXCEL

**MEDICATED
FOR RUMINANTS ONLY**

SHOW BUCK & DOE

FOR GROWING AND FINISHING MEAT GOATS FOR SHOW

For the prevention of coccidiosis caused by *Eimeria crandallis*, *Eimeria christenseni* and *Eimeria ninakohlykimovae*

ACTIVE DRUG INGREDIENTS

Monensin..... 20 g/ton

GUARANTEED ANALYSIS

| | | | |
|---|---------|------------------------------------|--------|
| CRUDE PROTEIN.....MIN | 16.00 % | SALT.....MIN | 0.70 % |
| This includes not more than 1.60% equivalent crude protein from non-protein nitrogen. | | SALT.....MAX | 1.00 % |
| CRUDE FAT.....MIN | 6.00 % | COBALT.....min.....PPM | 1.00 |
| CRUDE FIBER.....MAX | 17.00 % | COPPER.....min.....PPM | 26.00 |
| CALCIUM.....MIN | 0.90 % | COPPER.....max.....PPM | 30.00 |
| CALCIUM.....MAX | 1.20 % | MANGANESE...min.....PPM | 100.00 |
| PHOSPHORUS.....MIN | 0.35 % | ZINC.....min.....PPM | 105.00 |
| | | SELENIUM.....min.....PPM | 0.40 |
| | | VITAMIN ANOT LESS THAN IU/LB | 10,000 |
| | | VITAMIN ENOT LESS THAN IU/LB | 85 |

INGREDIENTS

ROUGHAGE PRODUCTS (25%), FORAGE PRODUCTS, GRAIN PRODUCTS, PROCESSED GRAIN BY-PRODUCTS, PLANT PROTEIN PRODUCTS, MOLASSES PRODUCTS, CALCIUM CARBONATE, ANIMAL PROTEIN PRODUCTS, AMMONIUM CHLORIDE, SALT, HYDROLYZED YEAST CELL WALL, DRIED ENTEROCOCCUS FAECIUM FERMENTATION PRODUCT, DRIED LACTOBACILLUS ACIDOPHILUS FERMENTATION PRODUCT, DRIED ASPERGILLUS ORYZAE, DRIED ASPERGILLUS NIGER FERMENTATION PRODUCT, DRIED TRICHODERMA LONGIBRACHIATUM FERMENTATION EXTRACT, DRIED BACILLUS SUBTILLUS FERMENTATION EXTRACT, ZINC SULFATE, L-LYSINE MONHYDROCHLORIDE, VEGETABLE OIL, NATURAL AND ARTIFICIAL FLAVORS, ZINC PROTEINATE, MANGANESE PROTEINATE, VITAMIN E SUPPLEMENT, SELENIUM YEAST, COPPER SULFATE, VITAMIN A SUPPLEMENT, COPPER PROTEINATE, VITAMIN D3 SUPPLEMENT, FERROUS SULFATE, COBALT PROTEINATE, ETHOXYQUIN (A PRESERVATIVE), ETHYLENEDIAMINE DIHYDRIDIDE, AND COBALT SULFATE.

CAUTION: 1. DO NOT FEED TO LACTATING GOATS. 2. DO NOT ALLOW HORSES OR OTHER EQUINES ACCESS TO FEED CONTAINING MONENSIN. INGESTION OF MONENSIN BY HORSES HAS BEEN FATAL. FEEDING UNDILUTED OR MIXING ERRORS RESULTING IN HIGH CONCENTRATIONS OF MONENSIN HAS BEEN FATAL IN CATTLE AND COULD BE FATAL IN GOATS. MONENSIN-MEDICATED FEED IS INTENDED FOR USE IN CATTLE AND IN GOATS. CONSUMPTION BY UNAPPROVED SPECIES MAY RESULT IN TOXIC REACTION.

FEEDING DIRECTIONS

FEEDING 1# OF EXCEL GOAT SUPPLIES 10 mg/lb OF MONENSIN. FEED CONTINUOUSLY AS THE SOLE RATION. PROVIDE FRESH WATER AT ALL TIMES. KEEP A GOOD QUALITY ROUGHAGE SUPPLY AVAILABLE WHEN NEEDED. FREE CHOICE MINERALS MAY BE OFFERED

081919 13440

ALLIED FEEDS INC.
208 HUTCHESON : CUERO, TEXAS 77954
NET WEIGHT 50# (22.72 kg)

01